

The lowest degree 0, 1-polynomial divisible by cyclotomic polynomial

A. Satyanarayana Reddy*

Abstract

Let n be an even positive integer with at most three distinct prime factors and let ζ_n be a primitive n -th root of unity. In this study, we made an attempt to find the lowest-degree 0, 1-polynomial $f(x)$ having at least three terms such that $f(\zeta_n)$ is a minimal vanishing sum of distinct n -th roots of unity.

1 Introduction and preliminaries

For a fixed positive integer n , let $U_n = \{k : 1 \leq k \leq n, \gcd(k, n) = 1\}$. If $|S|$ denotes the cardinality of the set S , then $|U_n| = \varphi(n)$, the well known *Euler-totient function*. Let $\zeta_n \in \mathbb{C}$ denote a primitive n -th root of unity, *i.e.*, $\zeta^n = 1$ and $\zeta^k \neq 1$ for $0 < k < n$. It is well known that the n -th cyclotomic polynomial, $\Phi_n(x) = \prod_{k \in U_n} (x - \zeta_n^k) \in \mathbb{Z}[x]$, is the minimal polynomial of ζ_n and $\deg(\Phi_n(x)) = \varphi(n)$. It is also known that $x^n - 1 = \prod_{d|n} \Phi_d(x)$ (here $a \mid b$ means ‘ a divides b ’). Now, using the property of minimal polynomials it follows that whenever $f(\zeta_n) = 0$ for some $f(x) \in \mathbb{Q}[x]$ then $\Phi_n(x)$ divides $f(x)$. Let $\Phi_n(x) = \sum_{k=0}^{\varphi(n)} a_{n,k} x^k$. Then it can be easily verified that $a_{n,i} = a_{n,\varphi(n)-i}$, where $0 \leq i \leq \varphi(n)$, $a_{n,0} = 1$ (for $n > 1$) and $a_{n,1} = -\sigma(n)$, where $\sigma(n) = \sum_{k \in U_n} \zeta_n^k$, is the sum of primitive n -th roots of unity. Before stating a result that gives the value of $a_{n,1}$, recall that a positive integer is said to be *square-free* if its decomposition into prime numbers/factors does not have any repeated factors.

Lemma 1.1 (Apostol [1]). *Let n be a positive integer. Then $\sigma(n) = \mu(n)$, where*

$$\mu(n) = \begin{cases} 0, & \text{if } n \text{ is not square-free,} \\ 1, & \text{if } n \text{ has even number of prime factors,} \\ -1, & \text{if } n \text{ has odd number of prime factors.} \end{cases}$$

For $f(x) \in \mathbb{Z}[x]$, we denote the set of coefficients of $f(x)$ by V_f and the set of exponents of $f(x)$ by E_f or $E_{f(x)}$. We define the *height* of the polynomial $f(x) \in \mathbb{Z}[x]$ to be the maximum absolute value of a coefficient of $f(x)$. Let $A(n)$ be the height of $\Phi_n(x)$. We say that a cyclotomic polynomial $\Phi_n(x)$ is *flat* if $A(n) = 1$. For $n < 105$, $\Phi_n(x)$ is flat. It was once conjectured that all cyclotomic polynomials are flat, however $A(105) = 2$. In fact, $A(n)$ is unbounded refer Emma Lehmer [7]. It is known that $\Phi_{pn}(x) = \Phi_n(x^p)$ whenever $p \mid n$ and $\Phi_{2n}(x) = \Phi_n(-x)$ whenever n is odd. Hence, if $\Phi_n(x)$ is a flat polynomial, then so is $\Phi_{pn}(x)$, where $p \mid n$ and $\Phi_{2n}(x)$ where n is odd.

*Department of Mathematics and Statistics, Indian Institute of Technology Kanpur, India 208016; (e-mail: satya@iitk.ac.in).

Let k be the number of distinct odd prime factors of n . For square-free n , this number k is called the *order* of the cyclotomic polynomial $\Phi_n(x)$. It is known that all cyclotomic polynomials of order 1 and order 2 are flat. Gennady Bachman [2] gave the first infinite family of flat cyclotomic polynomials of order three and this family was expanded by Kaplan [3]. In [4] Kaplan gave some flat polynomials of order four. It is unknown whether there are any flat cyclotomic polynomials of order greater than four.

Now, recall that an equation of the form $\zeta_n^{k_1} + \zeta_n^{k_2} + \cdots + \zeta_n^{k_l} = 0$ is called a *vanishing sum* of n -th roots of unity of weight l , where $k_1, k_2, \dots, k_l \in \{0, 1, 2, \dots, n-1\}$. The vanishing sum is said to be *minimal* if no proper sub-sums thereof can be zero.

For example, Let p be a prime divisor of n . Then $\zeta_n^{(n/p)k}$ for $1 \leq k \leq p-1$ are primitive p -th roots of unity and hence using Lemma 1.1, it follows that $1 + \sigma(p) = 0$ is a vanishing sum of n -th roots of unity and is in fact minimal as well.

Remark 1.2. *A vanishing sum can always be multiplied by a root of unity to get another vanishing sum. As such, the two vanishing sums are similar and in literature, it is said that the latter is obtained from the former by a rotation. Hence, the classification of minimal vanishing sums needs to be done only up to rotations (by roots of unity).*

We now state Corollary 3.2 of T.Y.Lam and K.H.Leung [6] that relates vanishing sums with square-free positive integers.

Lemma 1.3. *If $\alpha_1 + \alpha_2 + \cdots + \alpha_m = 0$ is a minimal vanishing sum of n -th roots of unity, then after a suitable rotation, we may assume that all α_i 's are n_0 -th roots of unity where n_0 is the largest square-free part of n .*

For example, if $n = p^k$ for some prime p , then $1 + \sigma(p) = 0$ is the only minimal vanishing sum of roots of unity up to rotation.

Lemma 1.4. *[T.Y.Lam and K.H.Leung [6]] Let $n = p^a q^b$, where p, q are distinct primes. Then, up to a rotation, the only minimal vanishing sums of n -th roots of unity are: $1 + \sigma(p) = 0$ and $1 + \sigma(q) = 0$.*

If n has 3 or more distinct prime factors, then T.Y.Lam and K.H.Leung [6] provided another minimal vanishing sum of n -th roots of unity, $\sigma(p_i) \cdot \sigma(p_j) + \sigma(p_k) = 0$ where p_i, p_j, p_k are distinct primes dividing n . Finding all minimal vanishing sums of n -th roots of unity when n has 3 or more distinct prime divisors seems difficult. In Section 2, we constructed another minimal vanishing sum of n -th roots of unity, when n is even and $\Phi_n(x)$ is flat. And we use this minimal vanishing sum for finding lowest 0,1 polynomial divisible by $\Phi_n(x)$. Note that in their paper, T.Y.Lam and K.H.Leung [6], allowed the repetition of n -th roots of unity in the vanishing sum. Gary Sivek [9] considered the vanishing sums of distinct n -th roots of unity. He found the number of terms in those vanishing sums. Unless specified otherwise, from now onwards, vanishing sum means vanishing sum of distinct n -th roots of unity.

John Steinberger [10] studied the lowest-degree polynomial with nonnegative coefficients divisible by $\Phi_n(x)$. He conjectured that the lowest degree monic polynomial with nonnegative coefficients divisible by $\Phi_n(x)$, $n > 1$, is $1 + x^{n/p} + \cdots + x^{(p-1)n/p}$ where p is the smallest prime dividing n . And proved this conjecture holds when n is even or when n is a prime power or when $2/p > 1/q_1 + \cdots + 1/q_k$ where $q_1, \dots, q_k$ are the other primes besides p dividing n . Consequently, if n is a odd prime power or when $2/p > 1/q_1 + \cdots + 1/q_k$ where $q_1, \dots, q_k$ are the other primes besides p dividing n and p is odd, then the lowest-degree 0,1- polynomial divisible by $\Phi_n(x)$ having at least three terms is $1 + x^{n/p} + \cdots + x^{(p-1)n/p}$.

Let $n = 2m$ for some $m \in \mathbb{Z}$. Then the lowest-degree 0, 1-polynomial divisible by $\Phi_n(x)$ is $x^{n/2} + 1$. Our interest in this paper, is to find the lowest-degree 0, 1-polynomial $f(x)$ with at least three terms such that $f(\zeta_{2m})$ is a minimal vanishing sum of $2m$ -th roots of unity. It is known (for example, see John P. Steinberger [11]) that the problem of finding the polynomials divisible by $\Phi_n(x)$ is equivalent to finding polynomials divisible by $\Phi_{n_0}(x)$, where n_0 is the maximum square-free factor of n . With this observation and Lemma 1.3, it is sufficient to consider square free n . Hence, in the remaining part of this paper, we consider $n = 2p_1 \dots p_k$, where $k \geq 1$ and $p_1 < p_2 < \dots < p_k$ are prime numbers.

Let us denote

$$G_n = \{f(x) | f(\zeta_n) \text{ is a minimal vanishing sum}\}.$$

Consequently, by our assumption of vanishing sum, if $f(x) \in G_n$, then $V_f \subseteq \{0, 1\}$, $\deg(f(x)) \leq n - 1$ and $\Phi_n(x) | f(x)$. Recall that a vanishing sum $f_1(\zeta_n) = 0$ is a rotation of another vanishing sum $f_2(\zeta_n) = 0$ if $f_1(\zeta_n) = \zeta_n^t f_2(\zeta_n)$ for some t . Now we define a relation $\mathcal{R}$ on G_n as $f_1(x) \mathcal{R} f_2(x)$ if and only if $f_1(\zeta_n) = 0$ is a rotation of $f_2(\zeta_n) = 0$. Clearly $\mathcal{R}$ is an equivalence relation on G_n . Let $H_n = G_n / \mathcal{R}$ be the set of equivalence classes of G_n under the equivalence relation $\mathcal{R}$. Without loss of generality we assume that the elements of H_n are the polynomials of least degree in the corresponding equivalence classes. Under this assumption $H_n \subseteq G_n$. Hence, if $f(x) \in H_n$, then $V_f \subseteq \{0, 1\}$, $f(\zeta_n)$ is a minimal vanishing sum, $\deg(f(x)) \leq n - 1$ and $f(0) = 1$. For example, let $n = p_0 p_1 \dots p_k$, $p_0 < p_1 < \dots < p_k$, $p_0 = 2$. Let $g_i(x) = 1 + \sum_{k=1}^{p_i-1} x^{\frac{kn}{p_i}} = 1 + x^{\frac{n}{p_i}} + x^{\frac{2n}{p_i}} + \dots + x^{\frac{(p_i-1)n}{p_i}}$. Then $g_i(x)$ is the polynomial corresponding to minimal vanishing sum $1 + \sigma(p_i) = 0$. Hence $g_i(x) \in G_n$. It is also easy to see that $g_i(x) \in H_n$ and $g_0(x) = x^{n/2} + 1$ is the only polynomial in H_n with two terms. With the definition of H_n , the problem of our interest is to find a lowest degree polynomial in H_n with at least three terms.

2 Minimal vanishing sum from flat cyclotomic polynomial

Let $n = p_0 p_1 p_2 \dots p_k$, where $k \geq 1$ and $p_0 = 2 < p_1 < p_2 < \dots < p_k$ are distinct primes. Suppose $\Phi_n(x)$ is flat. One can verify that

$$\Phi_n(x) = \begin{cases} x^{\varphi(n)} - x^{\varphi(n)-1} \pm \dots - x + 1, & \text{when } k \text{ odd,} \\ x^{\varphi(n)} + x^{\varphi(n)-1} - x^{\varphi(n)-p_1} \pm \dots - x^{p_1} + x + 1, & \text{when } k \text{ even.} \end{cases} \quad (1)$$

We can write $\Phi_n(x) = f_1(x) - f_2(x)$, where $f_1(x)$ and $f_2(x)$ are polynomials with positive coefficients. Observe this representation is unique as $E_{f_1} \cap E_{f_2} = \emptyset$. Now

$$\begin{aligned} \Phi(\zeta_n) &= 0 \\ \Rightarrow f_1(\zeta_n) - f_2(\zeta_n) &= 0 \\ \Rightarrow f_1(\zeta_n) + \zeta_n^{n/2} \cdot f_2(\zeta_n) &= 0 \\ \Rightarrow \Phi_n(x) | (f_1(x) + x^{n/2} f_2(x)). \end{aligned}$$

Let $\Phi_n^T(x) = f_1(x) + x^{n/2} f_2(x)$. Then $\Phi_n^T(x)$ is a 0, 1-polynomial. And from Equation (1) we have

$$\deg(\Phi_n^T(x)) = \begin{cases} \phi(n) - 1 + \frac{n}{2}, & \text{when } k \text{ is odd,} \\ \phi(n) - p_1 + \frac{n}{2}, & \text{when } k \text{ is even.} \end{cases} \quad (2)$$

Lemma 2.1. *Let $k \geq 1$ and $n = p_0 p_1 p_2 \dots p_k$, where $p_0 = 2 < p_1 < p_2 < \dots < p_k$ are distinct primes. Suppose $\Phi_n(x)$ is a flat polynomial. Then $\Phi_n^T(\zeta_n) = 0$ is a minimal vanishing sum of n -th roots of unity.*

Proof. Let us suppose $\Phi_n^T(\zeta_n) = 0$ is not a minimal vanishing sum of n -th roots of unity. Suppose $A(\zeta_n)$ is a proper sub-sum of $\Phi_n^T(\zeta_n)$ such that $A(\zeta_n) = 0$. Now we write $\Phi_n^T(x) = A(x) + B(x)$. Consequently $B(\zeta_n) = 0$. Without loss of generality assume that $A(x)$ does not contain the term $x^{\varphi(n)}$. We now write $A(x) = A_1(x) + A_2(x)$ such that the exponent of every term in $A_1(x) > n/2$ and that of $A_2(x) < n/2$. Then $\Phi_n(x)|A_2(x) - x^{-n/2}A_1(x)$ which is not possible since $\deg(A_2(x) - x^{-n/2}A_1(x)) < \varphi(n) = \deg(\Phi_n(x))$. $\square$

The following result is a direct consequence of Lemma 1.4. We are giving the proof for the sake of completeness.

Lemma 2.2. *Let p be any positive odd prime integer and $n = 2p$, then $H_n = \{g_0, g_1\}$. Further $g_1(x) = \Phi_n^T(x)$.*

Proof. First observe that $\Phi_{2p}(x) = \Phi_p(-x) = x^{p-1} - x^{p-2} + x^{p-3} - \dots - x + 1$. Hence $\Phi_n^T(x) = \sum_{i=0}^{p-1} x^{2i} = g_1(x)$. Let us suppose that $f(x) = x^{k_1} + x^{k_2} + \dots + x^{k_l} + 1 \in H_n$ where $n > k_1 > k_2 > \dots > k_l > 0$ and $f(x) \neq g_0(x)$. By definition of H_n , $\frac{n}{2} \notin E_f$. Now we write $f(x) = f_1(x) + f_2(x)$ in such a way that $\min(E_{f_1}) > \frac{n}{2}$ and $\max(E_{f_2}) < \frac{n}{2}$. Let $h(x) = f_2(x) - x^{\frac{n}{2}}f_1(x)$. Then degree of $\deg(h(x)) \leq \frac{n}{2} - 1 = p - 1 = \varphi(n)$. But $h(\zeta_n) = f_2(\zeta_n) - \zeta_n^{\frac{n}{2}}f_1(\zeta_n) = f(\zeta_n) = 0$. Hence $\Phi_n(x)|h(x)$ and $h(x) = \Phi_n(x)$. Thus $f(x) = \Phi_n^T(x) = g_1(x)$. $\square$

From the above lemma, the lowest degree polynomial in H_{2p} with at least three terms is $\Phi_{2p}^T(x)$. We now try to find a lowest degree polynomial in H_n with at least three terms, whenever $n = 2p_1p_2$, where $p_1 < p_2$ are odd primes. Since $g_1(x) \in H_n$, the degree of required polynomial is $\leq n - \frac{n}{p_1}$. In 1883, Migotti [8], showed that $\Phi_n(x)$ is a flat polynomial. Hence from Lemma 2.1, $\Phi_n^T(x) \in G_n$. In the following subsection, we show that $\Phi_n^T(x) \in H_n$.

2.1 $n = 2pq$

In this subsection, we suppose $n = 2pq$ where $p < q$ are odd primes. Lam and Heung [5] gave a nice expression for $\Phi_{pq}(x)$ as

$$\Phi_{pq}(x) = \left(\sum_{i=0}^r x^{ip} \right) \left(\sum_{j=0}^s x^{jq} \right) - \left(\sum_{i=r+1}^{q-1} x^{ip} \right) \left(\sum_{j=s+1}^{p-1} x^{jq} \right) x^{-pq}, \quad (3)$$

where r and s are positive integers such that $rp + sq = (p-1)(q-1) = \varphi(n)$, $0 \leq r \leq q-2$ and $0 \leq j \leq p-2$. Hence r and s have same parity.

Let us write Equation 3 as $\Phi_{pq}(x) = [A(x) + B(x)] - [C(x) + D(x)]$ such that $A(x) + B(x) = (\sum_{i=0}^r x^{ip})(\sum_{j=0}^s x^{jq})$ and $C(x) + D(x) = (\sum_{i=r+1}^{q-1} x^{ip})(\sum_{j=s+1}^{p-1} x^{jq})x^{-pq}$, where

$$\begin{aligned} E_A &= \{ip + jq | 0 \leq i \leq r, 0 \leq j \leq s \text{ and both } i \text{ and } j \text{ have same parity}\}, \\ E_B &= \{ip + jq | 0 \leq i \leq r, 0 \leq j \leq s \text{ and } i \text{ and } j \text{ are of different parity}\}. \end{aligned}$$

It is easy to see that $\max(E_A) = \varphi(n)$, $\min(E_A) = 0$, $\max(E_B) = \varphi(n) - p$ and $\min(E_B) = p$. Similarly we can define $C(x)$ and $D(x)$ such that $A(x)$, $C(x)$ are even functions and $B(x)$, $D(x)$ are odd functions.

With this we can write

$$\Phi_{2pq}(x) = [A(x) + D(x)] - [B(x) + C(x)] \text{ and } \Phi_{2pq}^T(x) = [x^{n/2}(B(x) + C(x))] + [A(x) + D(x)].$$

In order to prove $\Phi_n^T(x) \in H_n$ it is sufficient to prove that $\deg(\Phi_n^T(x)) \leq \deg(x^i \Phi_n^T(x))$ for all $i \in E_{\Phi_n^T}$. Hence the following result.

Theorem 2.3. Let $n = 2pq$ where $p, q (p < q)$ are odd primes. Then

1. $\deg(x^i \Phi_n^T(x)) > \deg(\Phi_n^T(x))$ for all $i \neq 0, \frac{n}{2} - p$.
2. $\deg(\Phi_n^T(x)) = \deg(x^{\frac{n}{2}-p} \Phi_n^T(x)) = \frac{n}{2} + \varphi(n) - p = n - 2p - q + 1$.

First we prove the following lemma.

Lemma 2.4. Let $n = 2pq$, where $p, q (p < q)$ are odd primes. Let $h \in E_{\Phi_n}$, $h < \varphi(n)$, then there exists $g \in E_{\Phi_n}$ such that $0 < g - h \leq p + q$.

Proof. If we prove the result for E_A and E_B , then it is also true of $E_{A(x)+D(x)}$ and $E_{B(x)+C(x)}$. Hence the result follows for E_{Φ_n} as $\deg(A(x)) - \deg(B(x)) = p$.

First we write $A(x) = A_1(x) + A_2(x)$ such that

$$\begin{aligned} E_{A_1} &= \{ip + jq | 0 \leq i \leq r, 0 \leq j \leq s \text{ and both } i \text{ and } j \text{ are even}\}, \\ E_{A_2} &= \{ip + jq | 0 \leq i \leq r, 0 \leq j \leq s \text{ and both } i \text{ and } j \text{ are odd}\}. \end{aligned}$$

Let $h \in E_A$. We will find $g \in E_A$ such that $g > h$ and $g - h \leq p + q$.

First consider the case when both r and s are even.

Suppose $h \in E_{A_1}$. Then $h = ip + jq$ where i, j are even, $0 \leq i \leq r$ and $0 \leq j \leq s$. Also $i = r, j = s$ is not possible since $h = rp + sq = \varphi(n)$.

Case1: $i \leq r - 2, j \leq s - 2 \Rightarrow g = (i + 1)p + (j + 1)q \in E_{A_2}$.

Case2: $i = r \text{ and } j \leq s - 2 \Rightarrow g = (r - 1)p + (j + 1)q \in E_{A_2}$.

Case3: $i \leq r - 2, j = s \Rightarrow g = (i + 2)p + sq \in E_{A_1}$.

Now suppose $h \in E_{A_2}$. Then $h = ip + jq$, i, j are odd, $0 \leq i \leq r - 1$ and $0 \leq j \leq s - 1$. In this case we can choose $g = (i + 1)p + (j + 1)q \in E_{A_1}$. With the same arguments we can prove that the difference between any two consecutive elements in E_A (with an abuse of language, assume there is an ordering in E_f) is at most $p + q$ when both r and s are odd.

Now we prove that the difference between any two consecutive elements in E_B is at most $p + q$. Before proceeding, recall $\varphi(n) - p$ is the largest element in E_B . Hence we suppose that $h < \varphi(n) - p$. As if $h = \varphi(n) - p$, then we choose $g = \varphi(n) \in E_A$ and $g - h < p + q$.

Let $h \in E_B$. Then $h = ip + jq$, where i, j have different parity and $0 \leq i \leq r$ and $0 \leq j \leq s$. Also by our assumption of h , $i = r - 1, j = s$ is not possible. We have the following cases.

Case1: i and r have same parity. Then j and s have different parity.

1. $i \leq r - 2, j \leq s - 1 \Rightarrow g = (i + 1)p + (j + 1)q$.
2. $i = r \text{ and } j \leq s - 1 \Rightarrow g = (r - 1)p + (j + 1)q$.

Case2: i and r have different parity. Then j and s have same parity.

1. $i \leq r - 1, j \leq s - 2 \Rightarrow g = (i + 1)p + (j + 1)q$.
2. $i < r - 1 \text{ and } j = s \Rightarrow g = (i + 2)p + sq$.

□

We now prove the Theorem 2.3.

Proof. Proof of Part 1: From the above lemma, the difference between any two consecutive terms in $E_{x^{\frac{n}{2}(C(x)+B(x))}}$ and $E_{A(x)+D(x)}$ is at most $p+q$. It is easy to see that the terms with the largest and the smallest exponent in $A(x)+D(x)$ belong to $A(x)$. Similar statement hold for $B(x)$ in $B(x)+C(x)$. Hence we have $\underbrace{\min(E_{x^{\frac{n}{2}(C(x)+B(x))}}) - \max(E_{A(x)+D(x)})}_{= \frac{n}{2} + p - \varphi(n)} = \underbrace{n - \max(E_{x^{\frac{n}{2}(C(x)+B(x))}})}_{= n - \max(E_{x^{\frac{n}{2}(C(x)+B(x))}})}$.

With these observations, the largest difference in the exponents of two consecutive terms in $\Phi_n^T(x)$ is $\frac{n}{2} + p - \varphi(n) = 2p + q - 1 (> p + q)$. Thus we conclude that $\deg(x^i \Phi_n^T(x)) > \deg(\Phi_n^T(x))$ for all $i \neq 0, \frac{n}{2} - p$.

Proof of Part 2: It is easy to see. $\square$

Now the degree of $\Phi_n^T(x)$ is $\frac{n}{2} + \varphi(n) - p = n - 2p - q + 1$, whereas the degree of $g_1(x) = n - \frac{n}{p} = n - 2q$, where $g_1(x)$ is the polynomial corresponding to minimal vanishing sum $1 + \sigma(p) = 0$. Now we conclude this work with the following conjecture and we feel that Lemma 2.5 given below, may be useful in proving this conjecture.

Conjecture: Let $n = 2pq$. Then the degree of lowest degree polynomial in H_n with at least three terms is $\min\{n - 2q, n - 2p - q + 1\}$.

Hence lowest degree polynomial(s) in H_n with at least three terms is (are)

$$\begin{cases} g_1(x), & \text{if } 2p < q + 1 \\ \Phi_n^T(x), & \text{if } 2p > q + 1 \\ \text{both } \Phi_n^T(x) \text{ and } g_1(x), & \text{if } 2p = q + 1. \end{cases}$$

For example, if $n = 30$, then $2p = q + 1$ and $\Phi_n^T(x) = x^{20} + x^{19} + x^{18} + x^8 + x^7 + x + 1$ whereas $g_1(x) = x^{20} + x^{10} + 1$.

Lemma 2.5. Let n be an even positive integer. Let $f(x) \in H_n$, where $|E_f| \geq 3$ and $\deg(f(x)) < \frac{n}{2} + \varphi(n)$. Then there exists $s \in E_f$ such that $\varphi(n) \leq s < \frac{n}{2}$.

Proof. From the definition of H_n , it is clear that $\frac{n}{2} \notin E_f$. So we can write $f(x) = f_1(x) + f_2(x)$ in such a way that $\min(E_{f_1}) > \frac{n}{2}$ and $\max(E_{f_2}) < \frac{n}{2}$. Then $\Phi_n(x) | f_2(x) - x^{-(n/2)} f_1(x)$. From the given hypothesis $\deg(x^{-(n/2)} f_1(x)) < \varphi(n)$, consequently $\max(E_{f_2}) \geq \varphi(n)$. Hence the result follows. $\square$

References

- [1] Tom M. Apostel, *Introduction to Analytic Number theory*, Springer-Verlag, New York, 1976.
- [2] Gennady Bachman, *Flat cyclotomic polynomials of order three*, Bull. London Math. Soc. 38 (2006) 5360.
- [3] Nathan Kaplan, *Flat cyclotomic polynomials of order three*, Journal of Number Theory, 127, 118-126 (2007).
- [4] Nathan Kaplan, *Flat cyclotomic polynomials of order four and higher*, Integers, 10, 357-363 (2010).
- [5] T.Y.Lam and K.H. Leung, *On the cyclotomic polynomial $\Phi_{pq}(x)$* , The American mathematical Monthly, Vol. 103, No. 7, 562-564 (Aug. - Sep., 1996).
- [6] T.Y.Lam and K.H. Leung, *On vanishing sums of roots of unity*, J. Algebra, 224, 91 -109 (2000).

- [7] Emma Lehmer, *On the magnitude of the coefficients of the cyclotomic polynomial*, Bull. Amer. Math. Soc., v. 42, pp. 389-392(1936).
- [8] A.Miggotti, *Zur Theorie der Kreisteilungsgleichung*, S.-B. der Math-Naturwiss. Classe der Kaiser Akad. der wiss., wien 87,7-14(1883).
- [9] Gary Sivek, *On vanishing sums of distinct roots of unity*, Integers, 10, 365-368 (2010).
- [10] John P. Steinberger, *The lowest-degree polynomial with nonnegative coefficients divisible by the n -th cyclotomic polynomial*, preprint.
- [11] John P. Steinberger, *Minimal Vanishing Sums of Roots of Unity with Large Coefficients*, Proc. London Math. Soc. (3) 97, 689-717, 2008.